

Comparative Study on the Reaction of Organophosphorus Reagents with *Moringa Oleifera* Vanillin. Synthesis of Phosphoranylidene-pyranone, Dioxaphospholane and Butenethione Derivatives as Antitumor Agents

M.A. Abd-El-Maksoud¹, S.A. Saleh², A.M. Abdallah², F.M. Soliman^{1*}

¹Department of and Organometalloid Chemistry, National Research Centre, 33-El-Buhouth St., P.O. 12622, Dokki, Giza, Egypt

²Horticultural Crops Technology Dept., National Research Centre, 33-El-Buhouth St., P.O. 12622, Dokki, Giza, Egypt

VANILLIN from *Moringa oleifera* plant was allowed to react with different nucleophilic organophosphorus reagents. Treatment of 4-hydroxy-3-methoxybenzaldehyde (vanillin) with the active phosphacumulene ylides (*N*-phenyliminovinylidene)- and (2-oxovinylidene)-triphenylphosphorane afforded the homocyclic phosphoranylidene-cyclobutylidenes. On the other hand, its reaction with the satabilized phosphoniumylide, acetylmethylenetriphenylphosphorane gave the corresponding butenone, which reacted with the active phosphacumulene to give finally the pyran derivative. Next, the phenyliminophenol derivative was obtained from the reaction of vanillin with the iminophosphorane. Moreover, dioxaphospholane oxide was isolated from the reaction of vanillin with phosphorus triamide. Finally, the butenethione was generated from the reaction of Japanese reagent with vanillin.

Keywords: *Moringa oleifera*, Vanillin, Phosphoranylidene, Heterocycles, Antitumors.

Introduction

Vanillin **1** is a naturally occurring substance. It is extracted from *Moringa oleifera* which is a multipurpose plant, utilized for medicinal applications such as cardiac and circulatory stimulants, antitumor, antiulcer, anti-inflammatory, antimicrobial, antihypertensive, antioxidant and antidiabetic agents [1-3]. Vanillin can be considered as one of the most important constituents of *Moringa* plant [4-6] and isolated also from tropical *Vanilla orchid*. It is also widely used in the food and beverage industry and is responsible for the characteristic vanilla flavor. This substance is also relevant to the synthesis of different agrochemicals, antifoaming and pharmaceutical products [7], such as antioxidants [8], antimicrobials [9-11], anti-mutagenic [12,13] and anti-carcinogenic agents [14]. Conversely, vanillin may also induce oxidative stress in yeast cells [15].

Experimental

Melting points were determined with an electrothermal digital melting point apparatus (Electro-Thermal Engineering Ltd., Essex, United Kingdom). The IR spectra were recorded in KBr

disks on a PyeUnicam SP 3300 and Shimadzu FT IR 8101 PC Infrared Spectrophotometers (PyeUnicam Ltd. Cambridge, England and Shimadzu, Tokyo, Japan, respectively). ¹H NMR spectra were obtained from a Jeol ECA 500 MHz NMR Spectrometer (Tokyo, Japan) using deuterated dimethylsulphoxide (d₆-DMSO) as a solvent and (TMS) as an internal reference at 500 MHz. Mass spectra (EI-MS) were obtained with ISQ (Single Quadrupole MS, Thermo Scientific). Elemental analyses agreed satisfactory with the calculated values. The using reported yields are of pure isolated materials obtained by column chromatography on silica gel 60 (Merck) and thin layer chromatography (TLC) which was performed on Merck Kiesel gel F254 precoated plates (Merck, Darmstadt, Germany). Solvents were dried/purified according to literature procedures.

Interaction of (N-phenyliminovinylidene) triphenylphosphorane (2a) and 4-hydroxy-3-methoxybenzaldehyde (1)

A solution of (N-phenyliminovinylidene) triphenylphosphorane (2a) [16] (0.377 g, 1 mmol)

*Corresponding author e-mail: solimanfma2@yahoo.com

DOI: 10.21608/EJCHEM.2018.2987.1253

©2017 National Information and Documentation Center (NIDOC)

in 20 mL of tetrahydrofuran, was added drop by drop with stirring at room temp, to a solution 4-hydroxy-3-methoxybenzaldehyde (1) (152g, 1 mmol) in 20 mL of tetrahydrofuran. The reaction mixture was stirred for 10 h during which the colour changed from yellow to dark brown (the progress of the reaction was monitored by TLC). THF was distilled off under reduced pressure and the residue was chromatographed on silica gel column using petroleum ether (60–80°C): ethyl acetate (30:70, v/v) as eluent. Product 5a was isolated along with triphenylphosphane oxide (m.p. and mixed m.p. 151°C).

2,4-Bis(phenylimino)-3-(triphenylphosphoranylidene)cyclobutylidene)methyl)-2-methoxyphenol (5a)

Yield 55%, yellow crystals, m.p. 270–272°C, IR (KBr, cm^{-1}): 3461 (OH), 1627 (2 C=N), 1557 (C=P), 1487, 1451 (P-Aryl). 1H NMR (500 MHz, DMSO, δ , ppm): 3.02 (s, 3 H, OCH₃), 6.28 (s, 1 H, OH, exchangeable with D₂O), 6.36–6.38 (d, 1 H, $^5J_{PH} = 10$ Hz, CH=P), 6.68–7.60 (m, 28H, Ar-H). ^{13}C NMR (125 MHz, d₆-DMSO, δ , ppm): 163.12, 163.03 (C=N), 153.48 (C=P), 118.37–138.72 (arom.-C). ^{31}P NMR (202.4 MHz, d₆-DMSO, δ , ppm): 29.79. MS(*m/z*, %): 628 [M]⁺, 6]. C₄₂H₃₃N₂O₂P (628.70).

Reaction of (2-oxovinylidene)triphenylphosphorane (2b) with 4-hydroxy-3-methoxybenzaldehyde (1)

A solution of (2-oxovinylidene)triphenylphosphorane (2b) [17] (302 g, 1 mmol) in 20 mL of dry toluene was added to a solution of 1 (0.152 g, 1 mmol) in 20 mL of dry toluene and refluxed for 10 h, during which the color was changed from colorless to yellow and precipitate is formed (the progress of the reaction was monitored by TLC). It was filtered and crystallized from ethyl acetate to give 5b and triphenylphosphane oxide.

2-(4-Hydroxy-3-methoxybenzylidene)-4-(triphenylphosphoranylidene)cyclobutane-1,3-dione (5b)

Yield 66%, colorless crystals, m.p. 309–310°C, IR (KBr, cm^{-1}): 3414 (OH), 1661 (br, 2 C=O), 1608 (C=C), 1504 (C=P), 1465, 1435 (P-aryl). 1H NMR (500 MHz, DMSO, δ , ppm): 3.02 (s, 3 H, OCH₃), 5.89 (s, 1 H, OH, exchangeable with D₂O), 6.79–7.92 (m, 19 H, Ar-H). ^{13}C NMR (125 MHz, d₆-DMSO, δ , ppm): 186.28 (2 C=O), 153.48 (C=P), 115.72–148.16 (arom.-C). ^{31}P NMR (202.4 MHz, d₆-DMSO, δ , ppm): 30.65. MS(*m/z*, %): 478 [M]⁺, 6]. C₃₀H₂₃O₄P (478.47).

Preparation of 4-hydroxy-3-methoxyphenyl)but-3-en-2-one (7)

A solution of 1-(triphenylphosphoranylidene)propan-2-one (6) (0.318 g) in 20 mL of dry toluene was added to a solution of 1 (0.152 g, 1 mmol) in 20 mL of dry toluene and refluxed for 15h, during which the colour was changed from colorless to yellow to dark brown, toluene was distilled off under reduced pressure and the residue was chromatographed on silica gel column using petroleum ether (60–80°C) and ethyl acetate. Product 7 was isolated along with triphenylphosphane oxide [18].

Interaction of (N-phenyliminovinylidene)triphenylphosphorane (2a) and 4-hydroxy-3-methoxyphenyl)but-3-en-2-one (7)

A mixture of (N-phenyliminovinylidene)triphenylphosphorane (2a) (0.377 g, 1 mmol) in 20 mL of dry THF, was added drop wise with stirring at room temperature, to a solution of 4-hydroxy-3-methoxyphenyl)but-3-en-2-one (7) (0.192 g, 1 mmol) in 20 mL of dry THF. The reaction mixture was stirred for 4 h during which the color was changed from yellow to red (the progress of the reaction was monitored by TLC). THF was distilled off and the residue was subjected to silica gel column chromatography using petroleum ether (60–80°C)/ethyl acetate as eluent (40: 60, v/v), to give compound 9. along with triphenylphosphane (m.p. and mixed m.p. 78°C).

2-Methoxy-4-(6-methyl-2-(phenylimino)-2H-pyran-4-yl)phenol (9)

Yield 45%, yellowish brown crystals, m.p. 120–122°C, IR (KBr, cm^{-1}): 3420 (OH). 1H NMR (500 MHz, DMSO, δ , ppm): 2.20 (s, 3 H, CH₃), 3.70 (s, 3 H, OCH₃), 5.07 (s, 1 H, =CH), 7.05–7.89 (m, 9 H, Ar-H), 12.52 (s, 1 H, OH, exchangeable with D₂O). MS (*m/z*, %): 307 [M⁺, 2], 230 [(M⁺-C₆H₅) 10]. C₁₉H₁₇NO₃ (307.34).

Reaction of (2-oxovinylidene)triphenylphosphorane (2b) with 4-hydroxy-3-methoxyphenyl)but-3-en-2-one (7)

A solution of (2-oxovinylidene)triphenylphosphorane (2b) (0.302 g, 1 mmol) in 20 mL of dry toluene was added to a solution of (0.192 mg, 1 mmol) in 20 mL of dry toluene and refluxed for 8h, during which the color was changed from yellow to brown (the progress of the reaction was monitored by TLC). Toluene was distilled off under reduced pressure and

the residue was subjected to silica gel column chromatography using petroleum ether (60–80°C)/ethyl acetate as an eluent (80: 20, v/v) to form compound **8b**.

4-(4-hydroxy-3-methoxyphenyl)-6-methyl-3-(triphenylphosphoranylidene)-3,6-dihydro-2H-pyran-2-one (8b)

Yield 67%, colorless crystals, m.p. 163-165°C, IR (KBr, cm^{-1}): 3425 (OH), 1680 (C=O), 1580 1480, 1437 (P-aryl). 1H NMR (500 MHz, DMSO, δ , ppm): 2.00 (s, 3 H, CH₃), 3.40 (s, 3 H, OCH₃), 5.82 (s, 1 H, OH, exchangeable with D₂O), 7.01-7.71 (m, 20 H, Ar-H). ^{31}P NMR (202.4 MHz, d_6 -DMSO, δ , ppm): 26.18. MS(m/z , %): 494 [M⁺], 6]. C₃₁H₂₇O₄P(494.52).

Interaction of (triphenylphosphoranylidene)aniline (12) and 4-hydroxy-3-methoxybenzaldehyde (1)

A solution of (triphenylphosphoranylidene)aniline (**12**) (0.353 g, 1 mmol) in 20 mL of dry toluene was added to a solution of **1** (0.152 g, 1 mmol) in 20 mL of dry toluene and refluxed for 15 h, during which the colour was changed from colorless to yellow and then dark brown (the progress of the reaction was monitored by TLC). Toluene was distilled off under reduced pressure and the residue was subjected to silica gel column chromatography using petroleum ether (60–80°C)/ethyl acetate as eluent (55 : 45, v/v) to form compound **13** and triphenylphosphane oxide.

2-Methoxy-4-((phenylimino)methyl)phenol (13)

Yield 45%, pale red crystals, m.p. 183-185°C, IR (KBr, cm^{-1}): 3417 (OH), 1590 (C=N). 1H NMR (500 MHz, DMSO, δ , ppm): 3.93 (s, 3 H, OCH₃), 5.94 (s, 1 H, OH, exchangeable with D₂O), 6.56 - 7.42 (m, 8 H, Ar-H), 7.46 (s, 1 H, CH=N). MS(m/z , %): 227 [M⁺], 6]. C₁₄H₁₃NO₂(227.26).

Interaction of hexamethyl phosphorus triamide (14) with 4-hydroxy-3-methoxybenzaldehyde (1)

A mixture of hexamethyl phosphorus triamide (**14**) (0.163 g, 1 mmol) in 20 mL of boiling toluene was added to a solution of **1** (0.152 g, 1 mmol) in 20 mL of dry toluene and refluxed for 3 h. Toluene is distilled off and the residue was crystallized from ethanol to form compound 18.

12-(dimethylamino)-3,5-bis(4-hydroxy-3-methoxyphenyl)-1,4,2-dioxaphospholane 2-oxide (18)

Yield 53%, colorless crystals, m.p. 111-113°C. 1H NMR (500 MHz, DMSO, δ , ppm): 2.63 (d,

6 H, 2 N-CH₃), 3.38 (s, 1 H, CH), 3.73 (s, 6 H, 2 OCH₃), 3.94 (s, 1 H, CH), 6.86- 7.38 (m, 6 H, Ar-H), 9.8 (s, 2 H, OH, exchangeable with D₂O). ^{31}P NMR (202.4 MHz, d_6 -DMSO, δ , ppm): 25.57. MS(m/z , %): 395 [M⁺], 10], 347 [M⁺-OCH₃+OH], 6], 299 [M⁺-2OCH₃+OH], 2]. C₁₈H₂₂NO₇P(395.34).

Reaction of (2,4-bis(phenylthio)-1,3,2,4-dithiadiphosphetane 2,4-disulfide) Japanese reagent (19) with 4-hydroxy-3-methoxyphenyl but-3-en-2-one (7)

A mixture of JR **19** (0.408 g, 1 mmol) in 20 mL of dry THF, was added drop wise with stirring at room temperature, to a solution of 4-hydroxy-3-methoxyphenylbut-3-en-2-one (**7**) (0.192 g, 1 mmol) in 20 mL of dry THF. The reaction mixture was stirred for 10 h (the progress of the reaction was monitored by TLC). THF was distilled off and the residue was subjected to silica gel column chromatography using petroleum ether (60–80°C)/ethyl acetate as eluent (75: 25, v/v) to form **20**.

(Z)-4-(4-Hydroxy-3-methoxyphenyl)but-3-ene-2-thione (20)

Yield 75%, colorless crystals, m.p. 235-237°C, IR (KBr, cm^{-1}): 3432 (OH), 1187 (C=S). 1H NMR (500 MHz, DMSO, δ , ppm): 2.45 (s, 3 H, CH₃), 3.4 (s, 3 H, OCH₃), 6.21 (s, 1 H, OH, exchangeable with D₂O), 6.89 (d, 1 H, J_{HH} = 5 Hz, CH), 7.28 (d, 1 H, J_{HH} = 5 Hz, CH), 7.46 - 7.78 (m, 3 H, Ar-H). ^{13}C NMR (125 MHz, d_6 -DMSO, δ , ppm): 24.23 (CH₃), 44.44 (OCH₃), 112 (CH), 120.56 -123.13 (arom.-C), 200.38 (C=S). C₁₁H₁₂O₂S(208.28).

Results and Discussion

Our interest in studying the reactions of organophosphorus reagents to synthesize pharmacologically interesting carbocyclic and heterocyclic systems containing phosphorus moieties [19-28] led us to examine the reactions of vanillin (**1**) with active phosphacumulenes, stabilized phosphonium ylides, phosphinimine, hexamethyl phosphorus triamide and Japanese reagent.


When vanillin (4-hydroxy-3-methoxybenzaldehyde) (**1**) was treated with two mole equivalent of active phosphacumulene ylide, namely, (*N*-phenylimino)vinylidene-triphenylphosphorane (**2a**) in THF at room temperature, the corresponding 2,4-bis(phenylimino)-3-(triphenylphosphoranylidene)cyclobutylidene)methyl-2-methoxyphenol (**5a**)

was obtained. Carrying out the reaction of vanillin 1 with (2-oxovinylidene) triphenylphosphorane (2b) in hot dry toluene afforded 2-(4-hydroxy-3-methoxybenzylidene)-4-(triphenylphosphoranylidene)cyclobutane-1,3-dione (5b). The phosphacumulenyliides can be described by the resonance structures 2A and 2B. So, initial nucleophilic attack by the carbanion center in the active phosphacumulene ylides (2a, b) furnishes the oxaphosphetane 3, as an intermediate [29-31]. Expulsion of triphenylphosphane oxide from 3 affords the unstable ketene 4 [32], which is followed immediately by [2+2]-cycloaddition to a second molecule of the active phosphacumulenes 2a,b giving the four-membered ring compounds 5a and b. Structural support for 5a was based upon correct elemental analysis and spectroscopic data. The IR spectrum of 5a (KBr, cm^{-1}) showed strong bands at 3461 (OH), 1627 (2 C=N), 1557 (C=P) [33], 1487, 1451 (P-Aryl) [34]. Moreover, 1H NMR (500 MHz, DMSO, δ , ppm) spectrum of 5a showed signals at : 3.02 (s, 3 H, OCH_3), 6.28 (s, 1 H, OH, exchangeable with D_2O), 6.36 - 6.38 (d,

1 H, CH=P), 6.68-7.60 (m, 28 H, Ar-H). There are signals at 163.12, 163.03 (C=N), 153.48 (C=P), 118.37-138.72 (arom.-C) in the ^{13}C NMR (125 MHz, d_6 -DMSO, δ , ppm) spectrum of 5a. A signal at δ 29.79 ppm was recorded in the ^{31}P NMR spectrum which fits with phosphorane in a four-membered ring [35-37] (Scheme 1).

The behavior of the stabilized phosphoniumylides, acetylmethylenetriphenylphosphorane 6 towards vanillin 1, afforded 4-hydroxy-3-methoxyphenylbut-3-en-2-one (7) together with triphenylphosphane oxide [38]. When the butenone 7 was treated with (N-phenyliminovinylidene)triphenylphosphorane (2a), the corresponding,

2-methoxy-4-(6-methyl-2-(phenylimino)-2H-pyran-4-yl)phenol (9) was produced. Formation of compound 9 is proposed to occur by [2+4]-cycloaddition of the ylide to compound 7 to give the phosphoranylidenepyran 8a. Since triphenylphosphane is a good leaving group, so



Scheme 1


Hoffmann degradation reaction occurs directly on compound 8a with the formation of the final product 9 and triphenylphosphane. On the other hand, reaction of (2-oxovinylidene) triphenylphosphorane (2b) with the butenone 7 afforded the stable 4-(4-Hydroxy-3-methoxyphenyl)-6-methyl-3-(triphenylphosphoranylidene)-3,6-dihydro-2*H*-pyran-2-one (8b). The most important features in the spectroscopic data of 8b is the appearance of a signal at δ 26.18 ppm and the mass spectrum showed an ion peak at m/z 494 $[M^+,6]$ (Scheme 2)

The reaction of phosphinimines are often analogous to those of phosphonium ylides. But in their activity iminophosphoranes are inferior to phosphinalkylenes [39-40]. We have found that 4-hydroxy-3-methoxybenzaldehyde (1) reacts with (triphenylphosphoranylidene)aniline (12) in boiling toluene to give 2-methoxy-4-((phenylimino)methyl)phenol (13), along with triphenylphosphane oxide. The mass spectrum of 13 showed a peak at m/z 227 $[(M)^+,6]$ (Scheme 3).

The reaction of hexamethyl phosphorustriamide (14) and the aldehyde 1 was investigated, too.

When the phosphorus reagent 14 was boiled with the aldehyde 1 in toluene, compound 2-(dimethylamino)-3,5-bis(4-hydroxy-3-methoxyphenyl)-1,4,2-dioxaphospholane 2-oxide (18) was obtained. As shown in scheme 4, formation of compound 18 is proposed to proceed by initial nucleophilic attack on the aldehyde 1 by hexamethylphosphorustriamide 14 to form the phosphonium species 15. Subsequent attack on another molecule of aldehyde 1 by the ylide 15 generates the intermediate 16 which can undergo ring closure to form 17. The last intermediate adds elements of water in the presence of unavoidable moisture to give the final stable product 18 together with dimethyl amine. The most important features in the spectroscopic data of the dioxaphospholane 18 is that it shows a signal at 25.57 ppm in P^{31} NMR spectrum which supports the dioxaphospholane structure [41]. The m/z was found at 395 $[M]^+$ in the mass spectrum of 18 (Scheme 4).

Chemistry of Japanese reagent (2,4-bis(phenylthio)-1,3,2,4-dithiadiphosphetane 2,4-disulfide) has been studied for many years [42]. This reagent 19a exists in equilibrium with the monomeric forms 19b, 19c, which can be incorporated with the substrate [43-45]. When the




Scheme 2


butene 7 was reacted with JR 19a, in dry *THF*, (4-hydroxy-3-methoxyphenyl)but-3-ene-2-thione (20) was formed. Only thionation reaction was happened and no addition reaction was observed between compound 7 and the monomer 19c (Scheme 5).

Conclusion


Vanillin 1 which can be extracted from the multipurpose plant *Moringa oleifera* was allowed to react with different organophosphorus reagents to prepare carbocyclic and heterocyclic compounds containing phosphorus which



Scheme 3



Scheme 4



Scheme 5

may be pharmacologically active. While vanillin reacts smoothly with the active (*N*-phenyliminovinylidene)-(2a) and (2-oxovinylidene)-triphenylphosphorane (2b), to give the four-membered ring phosphoranylidenecyclobutylidenes 5a,b, the stabilized phosphonium ylides (6) afforded the butenone (7), which reacted with the phosphacumulene (2a) to give the corresponding pyran derivatives (9). Moreover the difference in the nucleophilic character of 2a > 2b > 6 can be noticed in these reactions. These investigations afforded also the phenyliminophenol derivative (13) from the reaction of vanillin 1 with the iminophosphorane (12). Moreover the dioxaphospholane oxide derivative (18) was obtained from the reaction of phosphorus triamide 14 and the aldehyde vanillin 1. Finally, the butenethione 20 was produced from the reaction of the butenone 7 and Japanese reagent 19a. Only thionation reaction was observed in this investigation and no addition reaction between the butene 7 and the monomer of JR 19c occurred.

Acknowledgment

The authors would like to appreciate the Science and Technology Development Fund

(STDF), project no. 5979, which is gratefully acknowledged.

References

1. Eshak M.G., Hassanane M. M., Farag I.M., Shaffie N. M., Aboelfetoh M. A., Evaluation of protective and therapeutic role of *Moringa oleifera* leaf extract on CCL4-induced genotoxicity, hemotoxicity and hepatotoxicity in rats. *Intern. J. Pharm. Tech. Res.*, 7 (2), 392-415 (2015).
2. Abd El Baky H. H., El-Baroty G. S., Biological activity of the Egyptian *Moringa peregrina* seed oil, *Intern J Management Sciences Business Research*. 7(2), 98-108 (2013).
3. Asare G. A. I., Gyan B., Bugyei K., Adjei S. , Mahama R., Addo P., Otu-Nyarko L., Wiredu E. K., Nyarko A., Toxicity potentials of the nutraceutical *Moringa oleifera* at supra-supplementation levels. *J. Ethnopharmacol*, 139(1), 265-72 (2012) doi: 10.1016/j.jep.2011.11.009.
4. Abdelwanise F. M., Saleh S. A., Ezzo M. I., Helmy S. S. and Abodahab M. A., Response of *Moringa* plants to foliar application of Nitrogen and Potassium fertilizers. *Acta Horticulturae*, 1158, 187-194 (2017).
5. Hamed M. M., Abdalla A. M., Ghareeb M. A.

- and Saleh S. A. , Chemical constituents, in vitro antioxidant activity, oral acute toxicity and LD50 determination of *Moringa oleifera* leaves. *Inter. J. of Pharmacy and Pharmaceutical Sciences*, **9** (5), 240-247 (2017).
6. Saluja M. P., Kapil R. S., Popli S. P., Vanillin. *Indian J Chem., Sect. B*, **16**, 1044 (1978).
7. Walton N. J., Mayer M. J., Narbed A., Evaluation of antioxidant activity of vanillin by using multiple antioxidant assays. *Phytochemistry*, **63**, 505-515 (2003).
8. Tai A., Sawano T., Yazama, H.I., Analysis of the inhibition of food spoilage yeasts by vanillin, *Biochim Biophys Acta.*, **1810** (2), 170-7 (2011). doi: 10.1016/j.bbagen.2010.11.004.
9. Fitzgerald D.J., Stratford M., Narbad A., Synthesis, structural determination and antibacterial activity of compounds derived from vanillin and 4-aminoantipyrine, *Int J Food Microbiol.* **86** (1-2), 113-22 (2003).
10. Vaghasiya Y. K., Nair R., Soni M., Baluja S., Shanda S., Suppression of 6-TG-resistant mutations in V79 cells and recessive spot formations in mice by vanillin. *J. Serb. Chem. Soc.*, **69** (12), 991-998 (2004).
11. Rakchay S. S. , Suppakal P. , Jinkarn T. , Inhibitory effects of flavourings on mutagenesis induced by chemicals in bacteria. *As.J. Food Ag-In*, **2**, 138-147 (2009).
12. Imanishi H., Sasaki Y., Matsumoto K., Watanabe M., Ohta T., Shirasu Y., Tutikawa K., Suppressing effects of vanillin, cinnamaldehyde, and anisaldehyde on chromosome aberrations induced by X-rays in mice, *Mutation Res. Lett.*, **243** (2), 151-158 (1990). doi: 10.1016/0165-7992(90)90038.
13. Ohta T., Watanabe M., Watanabe K., Shirasu Y., Kada T., Vanillin causes the activation of Yap1 and mitochondrial fragmentation in *Saccharomyces cerevisiae*. *Food Chem Toxicol.* **24**(1), 51-4 (1986).
14. Ho K., Yazan L. S., Ismail N., Ismail M., Apoptosis and cell cycle arrest of human colorectal cancer cell line HT-29 induced by vanillin. *Cancer Epidemiol.* **33**(2), 155-60 (2009). doi: 10.1016/j.canep.2009.06.003
15. Nguyen T.T., Iwaki A., Ohya Y., Izawa S., Vanillin causes the activation of Yap1 and mitochondrial fragmentation in *Saccharomyces cerevisiae*, *J. Biosci Bioeng.*, **117**(1), 33-8 (2014). doi: 10.1016.
16. Bestmann H. J. and Schmid G., Sandmeier D., Synthesis of a Stable Alkylideneketenimine and an Alkylideneethioketene. *Angew Chem.*, **14**, 53-54 (1975). doi: 31239. 10.1002/anie.197500532.
17. Bestmann H. J., Sandmeier D., Simple Synthesis of Ketenylienetriphenylphosphorane and Its Thioanalogs, *Angew. Chem. Int. Edn.*, **14**, 634,(1975), *Chem. Abstr.*, **84**, 5070s (1976).
18. Hafez T. S., Henary M., Mahran M. R. H., The reaction of vanillin, Ortho-vanillin and piperonal with stabilized methylenetriphenylphosphoranes (Wittig reagents), *Phosphorus Sulfur Silicon Rel. Elem.*, **143**,: 33-44 (1998).
19. Shabana R., Maigali S.S., Essawy S.A., El-Hussieny M., Soliman F.M., Synthesis of thiaphosphetane, thiadiphosphetane and thiazadiphosphetidine derivatives from the reaction of phosphacumulenes, phosphallene and phosphinimin, *Egypt J Chem*, **50**, 59 (2007).
20. Maigali S.S., Abdel-Malek H. A., Soliman F. M., Reaction of camphorquinone derivatives with active phosphacumulenes and stabilised phosphonium ylides. *Egypt. J. Chem* **53** (2), 315-327 (2010).
21. Abd-El-Maksoud M. A., Maigali S. S., Soliman F. M., Facile Synthesis of Aziridine, Pyridine, Pyrolo[2,1-b]triazole Chromenones and Azaphosphinin Chromenones as Antitumor Agents, *J. Heterocycl. Chem.*, **51**, 1830-37 (2014).
22. Maigali S. S., Abd-El-Maksoud M. A., El-Hussieny M., Soliman F. M., Abdel-Aziz M. S., Shalaby S. M., Synthesis of antimicrobial agents from the reaction of aminoantipyrine, coumarin- and quinoline-carbaldehyde with phosphacumulene and phosphallene ylides *J. Chem. Res.*, **38**, 754-761 (2014).
23. Abd-El-Maksoud M. A., El-Hussieny M., Maigali S. S., Soliman F. M., Moharam M. E., The Reaction of Active Phosphonium Ylides with Aminothioxothiazolidinone, Thiazolamine. Synthesis and Antimicrobial Activity of Phosphanylidenes, Thiazolthione Pyrazoles and Cyclobutanes, *Res J Pharm Biol Chem Sci*, 1550-59 (2014).
24. Maigali S. S., Abd-El-Maksoud M. A., Soliman F. M., Moharam M. E., Synthesis of Pyrazoles by the [3 + 2] Cycloaddition of Diazo Compounds with Wittig Reagents as Antimicrobial Compounds, *J Heterocyclic Chem*, **52**, 834-40 (2015).
25. El-Hussieny M., Abd-El-Maksoud M. A., Maigali S. S., Soliman F. M., Soliman A. M., Chemistry of Phosphorus Ylides Part 42: Reaction of Dipyridyl Ethanedione with Phosphorus Reagents. Cytotoxic Activity of Phosphanylidene-Cyclobutane,

- Oxaphosphetane, and Pyridazinone Phosphorus, Sulfur, and Silicon, **190**, 1845–1856 (2015).
26. El-Hussieny M., Abd-El-Maksoud M. A., Maigali S. S., Soliman F. M., Chemistry of phosphorus ylides: Part 44. Reaction of 1-trimethylsilyl-1H-imidazole with phosphorus reagents. A convenient synthesis of phosphorus silyl imidazoles, *J. Chem. Res.* **40**, 265–68 (2016).
 27. Abd-El Maksoud M. A., Tawfik H. A., Maigali S. S., Soliman F. M., Moharam M. E., Dondeti M. F., Synthesis, antimicrobial and molecular docking evaluation of some heterocycles containing quinoline moiety, *Der Pharma Chemica*, **8** (13), 291-301(2016).
 28. Hashem A. I., EL-Hussieny, Abd-El-Maksoud M. M. A., Maigali S. S., Mansour Sh. T., Soliman F. M., Chemistry of phosphorus ylides: Part 45 synthesis of phosphoranylidene, thietane, azetidene and thiazinane derivatives as potent chemo preventative agents. *Phosphorus, Sulfur, and Silicon*, **193**, 1-9 (2018).
 29. Matthews C. N., Birum G. H., Carbophosphoranes, *Acc. Chem. Res.*, **2**, 373-339 (1969).
 30. Schlosser M., Piskala A., Tarchini C, Tuong H. B., Zum Ablauf der Wittig-Reaktion: zyklische oder ring offene Zwischenstufen? *Chimia*, **29**(8), 341-342 (1975).
 31. Vedejs E., Snoble K. A. J., Direct observation of oxaphosphetanes from typical Wittig reactions, *J. Am. Chem. Soc.*, **95**, 5778 (1973).
 32. Bestmann H. J., Schmid G., A New Synthesis of Keteniminylidene triphenylphosphoranes and Thioketenylidenetriphenylphosphorane. *Angew Chem Int Ed*, **13**, 273 (1974).
 33. Ramirez F., Madan O. P., Smith C. P., Reaction of trivalent phosphorus compounds with α,β -unsaturated ketones : Reaction of amino, alkyl and aryl phosphines with *trans*-dibenzoyl ethylene P nuclear magnetic resonance, *J. Org. Chem.*, **30**, 2284 (1965).
 34. Williams D. H., Fleming I., *Spectroscopic Methods In Organic Chemistry*, 4thed, McGraw-Hill Book Company, Maidenhead, Berkshire, UK, 55 (1987).
 35. Bestmann H. J., Schmid G., Sandmeier D., Kisielowski L., Dimere des Ketenyliden- und *N*-Phenyl-keteniminyliden-triphenylphosphorans. *Angew. Chem.*, **89**. 275 (1977); *Angew. Chem. Int. Ed. Engl.*, **16**, 268 (1977).
 36. Bestmann H. J., Zimmermann R., *Organic Phosphorus Compounds*, Vol. 3, Kosolopoff GM, Maier L (eds), New York: John Wiley and Sons, Inc. Wiley Chem. Abstr. 78 84461x (1973).
 37. Grim S.O., McFalarane W., Marks T.J., Phosphorus-31 chemical shifts of some air-sensitive phosphinemethylenes, *Chem Comm*, **1191** (1967). doi: 10.1039/C19670001191
 38. Wm. Johnson A., 'Ylid Chemistry in Organic Chemistry', A series of Monographs, Bolmquist AT, Academic Press, New York and London, (1966).
 39. Appel R., Hauss A., Zur Kenntnis des Imins, III. Phosphiniminium chloride und Triphenylphosphinimin, *Chem. Ber.*, **93**, 405–411 (1960).
 40. Horner L., Hoffmann H., Neuere Methoden der präparativen organischen Chemie II. Präparative und analytische Bedeutung tertiärer Phosphine und verwandter Verbindungen. Phosphor-organische Verbindungen VI, *Angew. Chem.*, **68**, 473 (1956).
 41. Pinkus A. G., Waldrep P. G., Ma S. Y., Structure of Products from Reactions of Phosphorus Pentachloride with Phenyl Salicylate and 2-Hydroxybenzophenone; Related Compounds. p^{31} N.M.R. and Chemical Studies *J. Heterocycl. Chem.*, **2**, 357-365 (1965).
 42. Cava M. P., Levinson M. I., Thionation reactions of lawesson's reagents, *Tetrahedron*, **41**, 5061-5087 (1985).
 43. Yoshifuji M., An, K. Toyota L.-D., Yasunami I. M., 2,4-Di-*t*-butyl-6-methoxyphenyldithioxophosphorane as a probe for the mechanistic studies of Lawesson's reagent. *Tetrahedr Lett.*, **35**, 4379-4382 (1994).
 44. Appel R., Knoch F., Kunze H., Das erste Organodithioxophosphoran. *Angew. Chem.*, **95**, 1008-1009 (1983).
 45. Bracher S., Cadogan J.I.G., Gosney I., Yaslak S., The generation and trapping of a monomeric aryldioxophosphorane ('metaphosphonate'), *J. Chem. Soc., Chem. Commun.*, 857-858(1983).

(Received 25/2/2018;
accepted 16/4/2018)

دراسة مقارنة عن تفاعل كواشف الفوسفور العضوية مع الفانيلين المستخرج من نبات المورينجا أوليفيرا
 تشييد مشتقات الفوسفور انيليديين بيرانون، و الداى أوكسافوسفولان و البيوتين ثيون ذات نشاط
 متوقع كمضادات للأورام

منصورة على عبد المقصود¹، ابو الفتوح محمد عبد الله²، سعيد عبد الحليم صالح³، فؤاد محمد سليمان⁴
¹ قسم الكيمياء العضوية الفلزية و العضوية شبه الفلزية - المركز القومي للبحوث - الجيزة - مصر.
² قسم تكنولوجيا الحاصلات البستانية - المركز القومي للبحوث - الجيزة - مصر.

نظرا للأهمية الفارماكولوجية لنبات المورينجا أوليفيرا لاستخداماته المتعددة فى المجالات الطبية و حيث أن مركب الفانيلين يعتبر من أهم مكونات هذا النبات. لذا فلقد تم مفاعلة مركب الفانيلين (٤-هيدروكسى-٣-ميثوكسى بنز الدهايد) مع كواشف الفوسفور النيكلوفيلية المترجمة النشطة و هى (ن-فينيل امينو فينيليديين)، (٢-أوكسوفينيليديين) ثلاثى فينيل الفوسفوران و التى أعطت مركبات متجانسة الحلقة و هى الفوسفور انيليديين سيكلوبيوتايلايديين. وعلى صعيد آخر وعند تفاعل الأيليدات الفوسفورية الثابتة مثل مشتق اسيتيل ميثيلين ثلاثى فينيل الفوسفوران مع مركب الفانيلين نتج عن هذا التفاعل مركب البيوتينون. و عند مفاعلة هذا البيوتينون مع الإيليدات الفوسفورية المترجمة تم فصل مشتقات البييران. و علاوة على ذلك تم الحصول على مركب أوكسيد الداى أوكسافوسفولون و ذلك من تفاعل الفانيلين مع كاشف ثلاثى اميد الفوسفور. و أخيرا تم فصل مركب البيوتين ثيون و ذلك من تفاعل كاشف جبانيز مع البيوتينون.